A-51

STATE KANSAS
COUNTY SEWARD
COMPANY Jomilson
FARM Light Estate "B"
WELL NO. 1
BLOCK
SURVEY
SEC. 11
T. 35s R. 32w
6342 TOTAL DEPTH
COMMENCED
COMPLETED
REMARKS C SW
ALTITUDE 2748
PRODUCTION 3000 oil + 47,500,000 gas 24hrs

CASING RECORD

SHOT	QUARTS	BETWEEN

FORM 186 - In stock and for sale by Ross-Martin Co., Tulsa

2100
2200
2300
Spls. Strt. 2400'
2400 wh.+ gray anhydrite with intercalated v.r.g. sh.
2500
Hollanberg 2538
g.bf, dse, mottled, selenitic dolo.
gray anhydrite
dolomite, some as above
tan suc. dolo
2600 tan to bf, suc. Δy dolo
gbf, suc. Δy dolo
tan to bf suc. Δy dolo.
good p.p. por. on tan to bf mottled suc. dolo.
2700 tan to bf, suc. Δy dolo
p.p. por. in cream suc. dolo.
g. dull op. Δ in part
2800 c-g, suc. dolo.
p.p. por. in gbf suc. dolo
Florence 2880
g.bf, f.gran, fos. dolomitic lst
2900 g, dull op. Δ, porosity in part
g.f. fos. fusulinid
g.f. gran. dolomitic lst
sl. Δy
t-g, d-f, fos.
c-g, d-f
3000 c-gbf, d-f, fos.
Wreford 2982
g. d-f
g.f.g. gran. Δy, dolomitic
3100 g.bf.g. shy, micac, silty
g.d-f, fos.
Stw Fusul. Zone 3090
c-tan, fusul. dolomitic
g. gran. dolomitic lst
Δy fusulinid fragment
g. d-f, fos.
N 3123
c-tan, Δy fairly ool. lst
3200 tan to g. gran. dolomitic lst
sl. Δy
dolomitic fusulinids
g. d-f, fos.
gbf, d-f, por. ooc. dolomitic lst
3300 g. d-f
c-tan, d-f fos.
g. dse
tan to gran. dolomitic lst
g. d-f
3400 g. d-f
g. gran. finely crenaceous lst
3500 g. micac, calc. sh.
g. d-f, fos.
g. d-f, d-f
Shly in part
3600 g. calc. sh. & marl
c-g. d-f, fos.
g. d-f, fos.
mottled pseudo-ool. in part
3700 g. fos. Δy
g. d-f, fos.
g. por. fusulinids
g. calc. sh.
g. f. fos. silty lst
3800 g. micac, sl. gritty sh
g. brn. f. fos. Δy
g. bf. f. fos.
g. gritty micac, calc. sst
g. d-f. fos. sl. Δy
g. calc. sh.
3900 g. micac. calc. sh.
c-g. d-f
g. bf. f
bf to brn. d-f
4000 highly nodular marl
g. calc, gritty micac, calc. sh.
bf calc
shly micac. sh
calc. silty micac sh
4100 g. gr. micac marl
mottled fos.
dk. g. micac sh
mottled fos. lst
4200 g. limey sh.
g. calc. micac. sh.
bf to brn. f. fos. shaley lst
4300 sh. calc. micac. st
silty lst
brn. f. gran. silty lst
med to dk. g. micac. sh.
bf. f. fos. shaley lst
4400 dk. g. micac. sh.
Hbnr Sh? 4354
med to dk. g. micac. sh.
dk. sh. micac. calc. sh
g. d-f
4500 g. d-f, fos. Δy
g. gran. d-f + dolomitic in streaks
4600 g. dse, Δy, sli. ooc. por.
g. fos.
Ls? 4547
g. d-f, sli. Δy, lst
4700 g. d-f, d-f
bf to brn. d-f
g. brn, d-f dk. brn. Δ
4800 g. d-f, Δy
g. d-f. Δy
g. bf. dse
ooc. lst
4900 g. d-f. Δy in part
dk. g. sh.
bf to brn. f. fos.
5000 g. bf. d-f, Δy
c-g. d-f. Δy
c-gbf, d-f, Δy
5100 g. d-f. fos. sl. ooc. lst
g. bf. d-f, Δy
g. brn. d-f, sli. Δy lst
bf to brn. d-f, Δy
5200 brn. f. Δy
bf to brn. d-f, Δy
dk. g. sh.
5300 g. brn, d-f
bf to brn. scat ooc. por.
d-f. Δy
g. d-f. fos. sli. Δy
c-g. d-f por. ooc. lst
dk. g. sh.
5400 tan to brn. d-f
f. gran.
dse. Δy
blk. sh.
g. d-f
5500 brn. d-f
g. d-f. dse
g. d-f. fos.
g. brn, dse Δy
g. ool. lst, sl. Δy
5600 g. d-f, fos. Δy
g. calc. sh.
g. brn. d-f, sli. Δy
5700
Δy
g. dse, sli. Δy
5800
d-f.
d-f, Δy
5628
Cherokee Sh. 5618
dk. g. calc. sh
5900 dk. brn. d-f. fos.
bf. sh.
lst interbedded with med to dk. g.
6000 med to dk. g. sh.
brn. dse Δy lst
g. d-f, sli. Δy
g. d-f. fos.
dk. g. dse Δy
blk. sh.
g. dse
brn. d-f
Atoka 5940
g. sh.
6100 ool. and in gray
spotty sand streaks
por. in bf f-med
calc sst.
t.o.s to very coarse grained
conglomeratic sh. spots of oil stn at top
gray, dull, coarse-textured
sh. with interbedded
g. bf. calc. gray s. f to
6200 g. flakey micac, limey sh
g. f. to c. xln. sandy fos. lst
phosphatic + glauconitic
brn. f. to c. xln. sdy fos. lst
gray flakey sl. sdy limey sh
bf to brn. dse to med. xln
brn. dse. Δy translu. Δ
bf to brn. pseudo-ool. lst
6300 interbedded with g. to c. xln. fos. lst
silty interbedded with g. sh.
Mississippian 6175
gray to bf. micro granular
to f. xln. fos. mottled pseudo-ool. interbedded with sh
as above
brn. to bf. dse to med. xln.
fos. lst.
"
6400
T.D. 6342
6500